Index of Authors

ABBERTON, M. T. *see* EASON, W. R.

ABRAHAM, E. M. and FASOULAS, A. C. Comparative efficiency of three selection methods in *Dactylis glomerata* L. and *Agropyron cristatum* L., 1

AHMAD, N. and WILMAN, D. The degradation of the cell walls of lucerne, Italian ryegrass and wheat straw when fed to cattle, sheep and rabbits, 337

ALBARRAN-PORTILLO, B. *see* ARRIAGA-JORDÁN, C. M.

ALLEN, E. J. *see* ALLISON, M. F.

ALLEN, E. J. *see* STALHAM, M. A.

ALLISON, M. F., FOWLER, J. H. and ALLEN, E. J. Effects of soil- and foliar-applied phosphorus fertilizers on the potato (*Solanum tuberosum*) crop, 1

ALLISON, M. F., FOWLER, J. H. and ALLEN, E. J. Factors affecting the magnesium nutrition of potatoes (*Solanum tubersosum*), 1

AL-SHOREPY, S. A. Genetic parameters for growth traits of a local breed of sheep in the United Arab Emirates, 365

ARCHIMEDE, H., DULORME, M., TOURNEBIZE, R., SAMINADIN, G., PERIACARPIN, F. and XANDE, A. The effects of *Gliricidia* supplementation on intake and digestion of a *Digitaria decumbens* hay by Black-belly sheep, 105

ARRIAGA-JORDÁN, C. M., FLORES-GALLEGOS, F. J., PEÑA-CARMONA, G., ALBARRAN-PORTILLO, B., GARCŒA-MARTŒNEZ, A., ESPINOZA-ORTEGA, A., GONZÁLEZ-ESQUIVEL, C. E. and CASTELÁN-ORTEGA, O. A. Participatory on-farm evaluation of the response to concentrate supplementation by cows in early lactation in smallholder peasant (*campesino*) dairy production systems in the highlands of central Mexico, 97

ASSAD, M. T. *see* DORRY, H. R.

BAKHEIT, B. R., ISMAIL, A. A., EL-SHIEMY, A. A. and SEDEK, F. S. Triple test cross analysis in four sesame crosses (*Sesamum indicum* L.), 1

BARRELL, G. K. *see* MAHGOUB, O.

BHATT, R. P., BISWAS, V. R. and KUMAR, N. Heterosis, combining ability and genetics for vitamin C, total soluble solids and yield in tomato (*Lycopersicon esculentum*) at 1700 m altitude, 71

BISWAS, P. K. and SALOKHE, V. M. Effects of planting date, intensity of tiller separation and plant density on the yield of transplanted rice, 279

BISWAS, V. R. *see* BHATT, R. P.

BONAMICO, N. C. *see* IBAÑEZ, M. A.

BRADBURY, N. J. *see* GABRIELLE, B.

BURMAN, U. *see* KATHJU, S.

BUTTER, N. L., DAWSON, J. M., WAKELIN, D. and BUTTERY, P. J. Effect of dietary condensed tannins on gastrointestinal nematodes, 1

BUTTERY, P. J. *see* BUTTER, N. L.

BYRNE, S. A. *see* WILMAN, D.

CABALLERO, M. *see* LORENZO, H.

CAO, B. *see* LU, X. G.

CARSON, A. F., IRWIN, D. and KILPATRICK, D. J. A comparison of Scottish Blackface and Cheviot ewes and five sire breeds in terms of lamb output at weaning in hill sheep systems, 1

CARSON, A. F., MOSS, B. W., DAWSON, L. E. R. and KILPATRICK, D. J. Effects of genotype and dietary forage to concentrate ratio during the finishing period on carcass characteristics and meat quality of lambs from hill sheep systems, 1

CASTELÁN-ORTEGA, O. A. *see* ARRIAGA-JORDÁN, C. M.

CHAMBERS, B. J. *see* WEBB, J.

CHEYGLINTED, S., RANAMUKHAARACHCHI, S. L. and SINGH, G. Assessment of the CERES-Rice model for rice production in the Central Plain of Thailand, 289

CHRISTAKIS, P. A. and FASOULAS, A. C. The recovery of recombinant inbreds outyielding the hybrid in tomato, 1

CULSHAW, C. M. *see* EASON, W. R.

DAWSON, J. M. *see* BUTTER, N. L.

DAWSON, L. E. R. *see* CARSON, A. F.

DHANOA, M. S. *see* EASON, W. R.

DHANOA, M. S. *see* SANDERSON, R.

DI RENZO, M. A. *see* IBAÑEZ, M. A.

DORRY, H. R. and ASSAD, M. T. Inheritance of leaf shape and its association with chlorosis in wheat infested by Russian wheat aphid (*Diuraphis noxia*), 1

DRAYCOTT, A. *see* RAHN, C.

DU, J.-Z. *see* HAN, X.-T.

DULORME, M. *see* ARCHIMEDE, H.

EASON, W. R., WEBB, K. J., MICHAELSON-YEATES, T. P. T., ABBERTON, M. T., GRIFFITH, G. W., CULSHAW, C. M., HOOKER, J. E. and DHANOA, M. S. Effect of genotype of *Trifolium repens* on mycorrhizal symbiosis with *Glomus mosseae*, 27

EKE-OKORO, O. N., OKEREKE, O. U. and OKEKE, J. E. Effect of stake sizes on some growth indices and yield of three cassava cultivars (*Manihot esculenta*), 1

EL-SHIEMY, A. A. *see* BAKHEIT, B. R.

ERSKINE, W. *see* SARKER, A.

ESPINOZA-ORTEGA, A. *see* ARRIAGA-JORDÁN, C. M.

FASOULAS, A. C. *see* ABRAHAM, E. M.

FASOULAS, A. C. *see* CHRISTAKIS, P. A.

FLORES-GALLEGOS, F. J. *see* ARRIAGA-JORDÁN, C. M.

FOULKES, M. J., SCOTT, R. K. and SYLVESTER-BRADLEY, R. The ability of wheat cultivars to withstand drought in UK conditions: resource capture, 1

FOWLER, J. H. *see* ALLISON, M. F.

GABRIELLE, B., RECOUS, S., TUCK, G., BRADBURY, N. J. and NICOLARDOT, B. Ability of the SUNDIAL model to

simulate the short-term dynamics of ^{15}N applied to winter wheat and oilseed rape.
GARCŒA-MARTŒNEZ, A. *see* ARRIAGA-JORDÁN, C. M.
GARG, B. K. *see* KATHJU, S.
GARWOOD, T. *see* WEBB, J.
GATES, F. *see* PELTONEN-SAINIO, P.
GHOSH, B. N. *see* MANNA, M. C.
GHOSH, P. K. *see* MANNA, M. C.
GOKANI, S. J. and THAKER, V. S. Accumulation of abscisic acid in cotton fibre and seed of normal and abnormal bolls,
GONZÁLEZ-ESQUIVEL, C. E. *see* ARRIAGA-JORDÁN, C. M.
GRIFFITH, G. W. *see* EASON, W. R.

HAN, X.-T., XUE, B., HU, L.-H. and DU, J.-Z. Effect of dietary protein degradability on net fluxes of free and peptide amino acids across the portal-drained viscera of steers,
HOLL, F. B. *see* TESFAYE, M.
HOOKER, J. E. *see* EASON, W. R.
HU, L.-H. *see* HAN, X.-T.
HUANG, Z. H. *see* LU, X. G.

IBAÑEZ, M. A., DI RENZO, M. A., SAMAME, S. S., BONAMICO, N. C. and POVERENE, M. M. Genotype–environment interaction of lovegrass forage yield in the semi-arid region of Argentina, 329
ILIADIS, C. Evaluation of six chickpea varieties for seed yield under autumn and spring sowing,
IRWIN, D. *see* CARSON, A. F.
ISMAIL, A. A. *see* BAKHEIT, B. R.

KATHJU, S., BURMAN, U. and GARG, B. K. Influence of nitrogen fertilization on water relations, photosynthesis, carbohydrate and nitrogen metabolism of diverse pearl millet genotypes under arid conditions, 307
KILPATRICK, D. J. *see* CARSON, A. F.
KIRKKARI, A-M. *see* PELTONEN-SAINIO, P.
KREUZER, M. *see* KÜLLING, D. R.
KRÖBER, T. F. *see* KÜLLING, D. R.
KÜLLING, D. R., MENZI, H., KRÖBER, T. F., NEFTEL, A., SUTTER, F., LISCHER, P. and KREUZER, M. Emissions of ammonia, nitrous oxide and methane from different types of manure during storage as affected by dietary protein content,
KUMAR, N. *see* BHATT, R. P.

LILLYWHITE, R. *see* RAHN, C.
LISCHER, P. *see* KÜLLING, D. R.
LORENZO, H., SIVERIO, J. M. and CABALLERO, M. Salinity and nitrogen fertilization and nitrogen metabolism in rose plants, 77
LOVELAND, P. J. *see* WEBB, J.
LU, X. G., YAO, K. M., YUAN, Q. H., CAO, B., MOU, T. M., HUANG, Z. H. and ZONG, X. M. Determination of fertility conversion indices for genic male-sterile rice in China,

MCALLAN, A. B. *see* SANDERSON, R.
MAHGOUB, O., BARRELL, G. K. and SYKES, A. R. Effect of gender, gonadectomy and oestradiol-17β on growth in lambs under grazing conditions, 351
MANNA, M. C., GHOSH, P. K., GHOSH, B. N. and SINGH, K. N. Comparative effectiveness of phosphate enriched compost and single superphosphate on yield, uptake of nutrients and soil quality under soybean–wheat rotation, 45
MEAD, A. *see* RAHN, C.
MENZI, H. *see* KÜLLING, D. R.
MERAH, O. Potential importance of water status traits for durum wheat improvement under Mediterranean conditions,
MICHAELSON-YEATES, T. P. T. *see* EASON, W. R.
MITCHELL, R. *see* WEBB, J.
MOSS, B. W. *see* CARSON, A. F.
MOU, T. M. *see* LU, X. G.
MUURINEN, S. *see* PELTONEN-SAINIO, P.

NEFTEL, A. *see* KÜLLING, D. R.
NICOLARDOT, B. *see* GABRIELLE, B.

OKEKE, J. E. *see* EKE-OKORO, O. N.
OKEREKE, O. U. *see* EKE-OKORO, O. N.
OWEN, H. LL. *see* WILMAN, D.

PELTONEN-SAINIO, P., MUURINEN, S., VILPPU, M., RAJALA, A., GATES, F. and KIRKKARI, A-M. Germination and grain vigour of naked oat in response to grain moisture at harvest,
PEÑA-CARMONA, G. *see* ARRIAGA-JORDÁN, C. M.
PERIACARPIN, F. *see* ARCHIMEDE, H.
PETERSEN, D. J. *see* TESFAYE, M.
POVERENE, M. M. *see* IBAÑEZ, M. A.

RAHN, C., MEAD, A., DRAYCOTT, A., LILLYWHITE, R. and SALO, T. A sensitivity analysis of the prediction of the nitrogen fertilizer requirement of cauliflower crops using the HRI WELL-N computer model, 55
RAJALA, A. *see* PELTONEN-SAINIO, P.
RANAMUKHAARACHCHI, S. L. *see* CHEYGLINTED, S.
RECOUS, S. *see* GABRIELLE, B.
ROWARTH, J. S. *see* WILLIAMS, P. H.
ROY, S. K., SHARMA, R. C. and TREHAN, S. P. Integrated nutrient management by using farmyard manure and fertilizers in potato–sunflower–paddy rice rotation in the Punjab,271
RUPA, T. R., SRIVASTAVA, S., SWARUP, A. and SINGH, D. Potassium supplying power of a Typic Ustochrept profile using quantity/intensity technique in a long-term fertilized plot,

SALO, T. A. *see* RAHN, C.
SALOKHE, V. M. *see* BISWAS, P. K.
SAMAME, S. S. *see* IBAÑEZ, M. A.
SAMINADIN, G. *see* ARCHIMEDE, H.
SANDERSON, R., DHANOA, M. S., THOMAS, C. and MCALLAN, A. B. Fish-meal supplementation of grass silage offered to young steers: Effects on growth, body composition and nutrient efficiency, 85
SARKER, A., SINGH, M. and ERSKINE, W. Efficiency of spatial methods in yield trials in lentil (*Lens culinaris* ssp. *culinaris*),
SCOTT, R. K. *see* FOULKES, M. J.
SEDEK, F. S. *see* BAKHEIT, B. R.
SHARMA, R. C. *see* ROY, S. K.
SINGH, D. *see* RUPA, T. R.
SINGH, G. *see* CHEYGLINTED, S.

SINGH, K. N. *see* MANNA, M. C.
SINGH, M. *see* SARKER, A.
SIVERIO, J. M. *see* LORENZO, H.
SRIVASTAVA, S. *see* RUPA, T. R.
STALHAM, M. A. and ALLEN, E. J. Effect of variety, irrigation regime and planting date on depth, rate, duration and density of root growth in the potato (*Solanum tuberosum*) crop, 251
SUTTER, F. *see* KÜLLING, D. R.
SWARUP, A. *see* RUPA, T. R.
SYKES, A. R. *see* MAHGOUB, O.
SYLVESTER-BRADLEY, R. *see* FOULKES, M. J.

TEJKLOVÁ, E. *see* WIESNER, I.
TESFAYE, M., PETERSEN, D. J. and HOLL, F. B. Nodulation responses of the tropical perennial clover, *Trifolium semipilosum*, inoculated with compatible and incompatible microsymbionts.
THAKER, V. S. *see* GOKANI, S. J.
THOMAS, C. *see* SANDERSON, R.
TOKATLIDIS, I. S. The effect of improved potential yield per plant on crop yield potential and optimum plant density in maize hybrids, 299
TOURNEBIZE, R. *see* ARCHIMEDE, H.
TREGURTHA, R. J. *see* WILLIAMS, P. H.
TREHAN, S. P. *see* ROY, S. K.
TUCK, G. *see* GABRIELLE, B.

VILPPU, M. *see* PELTONEN-SAINIO, P.

WAKELIN, D. *see* BUTTER, N. L.
WEBB, J., LOVELAND, P. J., CHAMBERS, B. J., MITCHELL, R. and GARWOOD, T. The impact of modern farming practices on soil fertility and quality in England and Wales.
WEBB, K. J. *see* EASON, W. R.
WIESNER, I., WIESNEROVÁ, D. and TEJKLOVÁ, E. Effect of anchor and core sequence in microsatellite primers on flax fingerprinting patterns, 37
WIESNEROVÁ, D. *see* WIESNER, I.
WILLIAMS, P. H., ROWARTH, J. S. and TREGURTHA, R. J. Uptake and residual value of ^{15}N-labelled fertilizer applied to first and second year grass seed crops in New Zealand, 17
WILMAN, D., BYRNE, S. A. and OWEN, H. LL. Long-term persistence of *Lolium perenne* and *Trifolium repens* in sown hill swards, 319
WILMAN, D. *see* AHMAD, N.

XANDE, A. *see* ARCHIMEDE, H.
XUE, B. *see* HAN, X.-T.

YAO, K. M. *see* LU, X. G.
YUAN, Q. H. *see* LU, X. G.

ZONG, X. M. *see* LU, X. G.

Index of Subjects

Abscisic acid
Accumulation of abscisic acid in cotton fibre and seed of normal and abnormal bolls,

Agropyron cristatum
Comparative efficiency of three selection methods in *Dactylis glomerata* L. and *Agropyron cristatum* L.,

Altitude
Heterosis, combining ability and genetics for vitamin C, total soluble solids and yield in tomato (*Lycopersicon esculentum*) at 1700 m altitude, 71

Amino acids
Effect of dietary protein degradability on net fluxes of free and peptide amino acids across the portal-drained viscera of steers,

Ammonia
Emissions of ammonia, nitrous oxide and methane from different types of manure during storage as affected by dietary protein content,

Aphid *see Diuraphis noxia*

Argentina
Genotype-environment interaction of lovegrass forage yield in the semi-arid region of Argentina, 329

Bolls
Accumulation of abscisic acid in cotton fibre and seed of normal and abnormal bolls,

Carbohydrate
Influence of nitrogen fertilization on water relations, photosynthesis, carbohydrate and nitrogen metabolism of diverse pearl millet genotypes under arid conditions, 307

Carcass
Effects of genotype and dietary forage to concentrate ratio during the finishing period on carcass characteristics and meat quality of lambs from hill sheep systems,

Cassava *see Manihot esculenta*

Cattle
The degradation of cell walls of *lucerne*, Italian ryegrass and wheat straw when fed to cattle, sheep and rabbits, 337

Cauliflower
A sensitivity analysis of the prediction of the nitrogen fertilizer requirement of cauliflower crops using the HRI WELL-N computer model, 55

Chickpea
Evaluation of six chickpea varieties for seed yield under autumn and spring sowing,

Concentrate
Effects of genotype and dietary forage to concentrate ratio during the finishing period on carcass characteristics and meat quality of lambs from hill sheep systems,

Conversion
Determination of fertility conversion indices for genic male-sterile rice in China,

Cotton
Accumulation of abscisic acid in cotton fibre and seed of normal and abnormal bolls,

Cows
Participatory on-farm evaluation of the response to concentrate supplementation by cows in early lactation in smallholder peasant (*campesino*) dairy production systems in the highlands of central Mexico, 97

Dactylis glomerata
Comparative efficiency of three selection methods in *Dactylis glomerata* L. and *Agropyron cristatum* L.,

Degradability
Effect of dietary protein degradability on net fluxes of free and peptide amino acids across the portal-drained viscera of steers,

Digitaria decumbens
The effects of *Gliricidia* supplementation on intake and digestion of a *Digitaria decumbens* hay by Black-belly sheep, 105

Diuraphis noxia
Inheritance of leaf shape and its association with chlorosis in wheat infested by Russian wheat aphid (*Diuraphis noxia*),

Drought
The ability of wheat cultivars to withstand drought in UK conditions: resource capture, 1

Durum wheat
Potential importance of water status traits for durum wheat improvement under Mediterranean conditions,

Ewes
A comparison of Scottish Blackface and Cheviot ewes and five sire breeds in terms of lamb output at weaning in hill sheep systems,

Farming practices
The impact of modern farming practices on soil fertility and quality in England and Wales,

Fertility
Determination of fertility conversion indices for genic male-sterile rice in China,

Fertilizer
Uptake and residual value of ^{15}N-labelled fertilizer applied to first and second year grass seed crops in New Zealand, 17

A sensitivity analysis of the prediction of the nitrogen fertilizer requirement of cauliflower crops using the HRI WELL-N computer model, 55

Integrated nutrient management by using farmyard manure and fertilizers in potato-sunflower-paddy rice rotation in the Punjab, 271

Fish-meal
Fish-meal supplementation of grass silage offered to young steers: Effects on growth, body composition and nutrient efficiency, 85

Flax
Effect of anchor and core sequence in microsatellite primers on flax fingerprinting patterns, 37
Foliar
Effects of soil- and foliar-applied phosphorus fertilizers on the potato (*Solanum tuberosum*) crop,
Forage
Effects of genotype and dietary forage to concentrate ratio during the finishing period on carcass characteristics and meat quality of lambs from hill sheep systems,
Gastrointestinal
Effect of dietary condensed tannins on gastrointestinal nematodes,
Gender
Effect of gender, gonadectomy and oestradiol-17β on growth in lambs under grazing conditions, 351
Genetic Parameters
Genetic parameters for growth traits of a local breed of sheep in the United Arab Emirates, 365
Genotype
Effect of genotype of *Trifolium repens* on mycorrhizal symbiosis with *Glomus mosseae*, 27
Germination
Germination and grain vigour of naked oat in response to grain moisture at harvest,
Gliricidia
The effects of *Gliricidia* supplementation on intake and digestion of a *Digitaria decumbens* hay by Black-belly sheep, 105
Glomus mosseae
Effect of genotype of *Trifolium repens* on mycorrhizal symbiosis with *Glomus mosseae*, 27
Gonadectomy
Effect of gender, gonadectomy and oestradiol-17β on growth in lambs under grazing conditions, 351
Grain vigour
Germination and grain vigour of naked oat in response to grain moisture at harvest,
Grass
Uptake and residual value of ^{15}N-labelled fertilizer applied to first and second year grass seed crops in New Zealand, 17
Fish-meal supplementation of grass silage offered to young steers: Effects on growth, body composition and nutrient efficiency, 85
Growth
Effect of stake sizes on some growth indices and yield of three cassava cultivars (*Manihot esculenta*),
Hay
The effects of *Gliricidia* supplementation on intake and digestion of a *Digitaria decumbens* hay by Black-belly sheep, 105
Indices
Determination of fertility conversion indices for genic male-sterile rice in China,
Irrigation
Effect of variety, irrigation regime and planting date on depth, rate, duration and density of root growth in the potato (*Solanum tubersosum*) crop, 251
Italian ryegrass
The degradation of cell walls of *lucerne*, Italian ryegrass and wheat straw when fed to cattle, sheep and rabbits, 337
Lactation
Participatory on-farm evaluation of the response to concentrate supplementation by cows in early lactation in smallholder peasant (*campesino*) dairy production systems in the highlands of central Mexico, 97
Lambs
Effects of genotype and dietary forage to concentrate ratio during the finishing period on carcass characteristics and meat quality of lambs from hill sheep systems,
A comparison of Scottish Blackface and Cheviot ewes and five sire breeds in terms of lamb output at weaning in hill sheep systems,
Effect of gender, gonadectomy and oestradiol-17β on growth in lambs under grazing conditions, 351
Lens culinaris
Efficiency of spatial methods in yield trials in lentil (*Lens culinaris* ssp. *culinaris*),
Lentil *see Lens culinaris*
Lolium perenne
Long-term persistence of *Lolium perenne* and *Trifolium repens* in sown hill swards, 319
Lovegrass
Genotype-environment interaction of lovegrass forage yield in the semi-arid region of Argentina, 329
Lucerne
The degradation of cell walls of *lucerne*, Italian ryegrass and wheat straw when fed to cattle, sheep and rabbits, 337
Lycopersicon esculentum
Heterosis, combining ability and genetics for vitamin C, total soluble solids and yield in tomato (*Lycopersicon esculentum*) at 1700 m altitude, 71
The recovery of recombinant inbreds outyielding the hybrid in tomato,
Magnesium
Factors affecting the magnesium nutrition of potatoes (*Solanum tubersosum*),
Maize
The effect of improved potential yield per plant on crop yield potential and optimum plant density in maize hybrids, 299
Manihot esculenta
Effect of stake sizes on some growth indices and yield of three cassava cultivars (*Manihot esculenta*),
Manure
Emissions of ammonia, nitrous oxide and methane from different types of manure during storage as affected by dietary protein content,
Integrated nutrient management by using farmyard manure and fertilizers in potato-sunflower-paddy rice rotation in the Punjab, 271
Meat
Effects of genotype and dietary forage to concentrate ratio during the finishing period on carcass characteristics and meat quality of lambs from hill sheep systems,
Mediterranean
Potential importance of water status traits for durum wheat improvement under Mediterranean conditions,
Methane
Emissions of ammonia, nitrous oxide and methane from different types of manure during storage as affected by dietary protein content,

Microsatellite
Effect of anchor and core sequence in microsatellite primers on flax fingerprinting patterns, 37
Microsymbionts
Nodulation responses of the tropical perennial clover, *Trifolium semipilosum*, inoculated with compatible and incompatible microsymbionts,
Model
Ability of the SUNDIAL model to simulate the short-term dynamics of ^{15}N applied to winter wheat and oilseed rape,
Moisture
Germination and grain vigour of naked oat in response to grain moisture at harvest,
Mycorrhizal
Effect of genotype of *Trifolium repens* on mycorrhizal symbiosis with *Glomus mosseae*, 27
Nematodes
Effect of dietary condensed tannins on gastrointestinal nematodes,
Nitrogen
A sensitivity analysis of the prediction of the nitrogen fertilizer requirement of cauliflower crops using the HRI WELL-N computer model, 55
Salinity and nitrogen fertilization and nitrogen metabolism in rose plants, 77
Influence of nitrogen fertilization on water relations, photosynthesis, carbohydrate and nitrogen metabolism of diverse pearl millet genotypes under arid conditions, 307
Nitrous oxide
Emissions of ammonia, nitrous oxide and methane from different types of manure during storage as affected by dietary protein content,
Nodulation
Nodulation responses of the tropical perennial clover, *Trifolium semipilosum*, inoculated with compatible and incompatible microsymbionts,
Oat
Germination and grain vigour of naked oat in response to grain moisture at harvest,
Oestradiol-17*β*
Effect of gender, gonadectomy and oestradiol-17β on growth in lambs under grazing conditions, 351
Oilseed rape
Ability of the SUNDIAL model to simulate the short-term dynamics of ^{15}N applied to winter wheat and oilseed rape,
Paddy-rice
Integrated nutrient management by using farmyard manure and fertilizers in potato-sunflower-paddy rice rotation in the Punjab, 271
Pearl millet
Influence of nitrogen fertilization on water relations, photosynthesis, carbohydrate and nitrogen metabolism of diverse pearl millet genotypes under arid conditions, 307
Perennial clover *see Trifolium semipilosum*
Phosphate
Comparative effectiveness of phosphate enriched compost and single superphosphate on yield, uptake of nutrients and soil quality under soybean–wheat rotation, 45
Phosphorus
Effects of soil- and foliar-applied phosphorus fertilizers on the potato (*Solanum tuberosum*) crop,
Photosynthesis
Influence of nitrogen fertilization on water relations, photosynthesis, carbohydrate and nitrogen metabolism of diverse pearl millet genotypes under arid conditions, 307
Potassium
Potassium supplying power of a Typic Ustochrept profile using quantity/intensity technique in a long-term fertilized plot,
Potato *see Solanum tubersosum*
Protein
Emissions of ammonia, nitrous oxide and methane from different types of manure during storage as affected by dietary protein content,
Effect of dietary protein degradability on net fluxes of free and peptide amino acids across the portal-drained viscera of steers,
Punjab
Integrated nutrient management by using farmyard manure and fertilizers in potato-sunflower-paddy rice rotation in the Punjab, 271
Rabbits
The degradation of cell walls of *lucerne*, Italian ryegrass and wheat straw when fed to cattle, sheep and rabbits, 337
Rice
Effects of planting date, intensity of tiller separation and plant density on the yield of transplanted rice, 279
Assessment of the CERES-Rice model for rice production in the Central Plain of Thailand, 289
Determination of fertility conversion indices for genic male-sterile rice in China,
Root
Effect of variety, irrigation regime and planting date on depth, rate, duration and density of root growth in the potato (*Solanum tubersosum*) crop, 251
Rose
Salinity and nitrogen fertilization and nitrogen metabolism in rose plants, 77
Seed
Uptake and residual value of ^{15}N-labelled fertilizer applied to first and second year grass seed crops in New Zealand, 17
Sesame *see Sesamum indicum*
Sesamum indicum
Triple test cross analysis in four sesame crosses (*Sesamum indicum* L.),
Sheep
The degradation of cell walls of *lucerne*, Italian ryegrass and wheat straw when fed to cattle, sheep and rabbits, 337
Genetic parameters for growth traits of a local breed of sheep in the United Arab Emirates, 365
Silage
Fish-meal supplementation of grass silage offered to young steers: Effects on growth, body composition and nutrient efficiency, 85
Sire
A comparison of Scottish Blackface and Cheviot ewes and five sire breeds in terms of lamb output at weaning in hill sheep systems,

Soil
Comparative effectiveness of phosphate enriched compost and single superphosphate on yield, uptake of nutrients and soil quality under soybean–wheat rotation, 45
Effects of soil- and foliar-applied phosphorus fertilizers on the potato (*Solanum tuberosum*) crop,
Soil fertility
The impact of modern farming practices on soil fertility and quality in England and Wales,
Solanum tubersosum
Effect of variety, irrigation regime and planting date on depth, rate, duration and density of root growth in the potato (*Solanum tubersosum*) crop, 251
Integrated nutrient management by using farmyard manure and fertilizers in potato-sunflower-paddy rice rotation in the Punjab, 271
Effects of soil- and foliar-applied phosphorus fertilizers on the potato (*Solanum tuberosum*) crop,
Factors affecting the magnesium nutrition of potatoes (*Solanum tubersosum*),
Soybean
Comparative effectiveness of phosphate enriched compost and single superphosphate on yield, uptake of nutrients and soil quality under soybean–wheat rotation, 45
Spatial methods
Efficiency of spatial methods in yield trials in lentil (*Lens culinaris* ssp. *culinaris*),
Steers
Fish-meal supplementation of grass silage offered to young steers: Effects on growth, body composition and nutrient efficiency, 85
Effect of dietary protein degradability on net fluxes of free and peptide amino acids across the portal-drained viscera of steers,
Sunflower
Integrated nutrient management by using farmyard manure and fertilizers in potato-sunflower-paddy rice rotation in the Punjab, 271
Superphosphate
Comparative effectiveness of phosphate enriched compost and single superphosphate on yield, uptake of nutrients and soil quality under soybean–wheat rotation, 45
Tannins
Effect of dietary condensed tannins on gastrointestinal nematodes,
Thailand
Assessment of the CERES-Rice model for rice production in the Central Plain of Thailand, 289
Tiller
Effects of planting date, intensity of tiller separation and plant density on the yield of transplanted rice, 279
Tomato *see Lycopersicon esculentum*
Trifolium repens
Effect of genotype of *Trifolium repens* on mycorrhizal symbiosis with *Glomus mosseae*, 27
Long-term persistence of *Lolium perenne* and *Trifolium repens* in sown hill swards, 319
Trifolium semipilosum
Nodulation responses of the tropical perennial clover, *Trifolium semipilosum*, inoculated with compatible and incompatible microsymbionts,
United Arab Emirates
Genetic parameters for growth traits of a local breed of sheep in the United Arab Emirates, 365
Water
Potential importance of water status traits for durum wheat improvement under Mediterranean conditions,
Influence of nitrogen fertilization on water relations, photosynthesis, carbohydrate and nitrogen metabolism of diverse pearl millet genotypes under arid conditions, 307
Wheat
The ability of wheat cultivars to withstand drought in UK conditions: resource capture, 1
Comparative effectiveness of phosphate enriched compost and single superphosphate on yield, uptake of nutrients and soil quality under soybean–wheat rotation, 45
Ability of the SUNDIAL model to simulate the short-term dynamics of ^{15}N applied to winter wheat and oilseed rape,
Inheritance of leaf shape and its association with chlorosis in wheat infested by Russian wheat aphid (*Diuraphis noxia*),
Wheat straw
The degradation of cell walls of *lucerne*, Italian ryegrass and wheat straw when fed to cattle, sheep and rabbits, 337
Yield
Heterosis, combining ability and genetics for vitamin C, total soluble solids and yield in tomato (*Lycopersicon esculentum*) at 1700 m altitude, 71
Effect of stake sizes on some growth indices and yield of three cassava cultivars (*Manihot esculenta*),
Efficiency of spatial methods in yield trials in lentil (*Lens culinaris* ssp. *culinaris*),
Evaluation of six chickpea varieties for seed yield under autumn and spring sowing,